

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



7-84M
op. 1

LIBRARY
RECEIVED
1943

U. S. Department of Agriculture



VICTORY GARDENS

MISCELLANEOUS
PUBLICATION
NO. 483 *rev.*
Jan. 1943

U. S. DEPARTMENT OF AGRICULTURE

VEGETABLES that are common and easily grown can furnish a good share of the energy, vitamins, and minerals that all of us need every day. During periods of national emergency normal commercial vegetable supplies may be too short to meet unusual demands, or otherwise not within reach of everyone. It then becomes a public duty, as well as a private necessity, for everyone who is able to grow at least a part of the vegetables needed for his family.

Beginners who are able to make a vegetable garden are more likely to succeed if they confine their efforts to a few staple and productive kinds of crops, such as those mentioned in the following pages.

In the brief space of a publication of this character only the bare outlines of simple vegetable gardens can be presented. This publication is intended to give only general information that will introduce the inexperienced prospective gardener to the subject and enable him to make a beginning. It gives elementary information on what to grow, how to prepare and fertilize the soil, how and when to plant, how to care for the plants, and how to utilize the crop. As an aid to success, "Victory gardeners" should consult their State agricultural colleges, county agents, vocational agricultural teachers, local experienced gardeners, and should study the publications on gardening and storage more closely adapted to their local conditions, prepared by their own State agricultural experiment stations and extension services. The United States Department of Agriculture also publishes several more comprehensive bulletins on the subject. The following can be obtained free from the Office of Information, United States Department of Agriculture, Washington, D. C.:

Leaflet 203, Disease-Resistant Varieties of Vegetables for the Home Garden.

Farmers' Bulletin 1044, The City Home Garden.

Farmers' Bulletin 1371, Diseases and Insects of Garden Vegetables.

Farmers' Bulletin 1673, The Farm Garden.

Farmers' Bulletin 1743, Hotbeds and Coldframes.

The following can be purchased for 5 cents from the Superintendent of Documents, Government Printing Office, Washington, D. C.:

Department Bulletin 1427, Dry-Land Gardening at the Northern Great Plains Field Station, Mandan, N. Dak.

Unusual demands are being made on domestic supplies of vegetable seeds and fertilizers. Do not waste them.


Tools are scarce. Take good care of them.



VICTORY GARDENS

By VICTOR R. BOSWELL

Principal horticulturist and assistant head of division, Division of Fruit and Vegetable Crops and Diseases, Bureau of Plant Industry, Agricultural Research Administration



VICTORY GARDENS ARE ESSENTIAL

IN TIMES OF WAR the country's food supply for civilians may be smaller even though total production is greatly increased. The burdens of World War II are already creating serious food-production, distribution, and preservation problems. Labor and machinery shortages interfere with production; overloaded railroads and restricted motor transport interfere with distribution; and inadequate supplies of labor, steel, and tin demand that civilians depend less on foods canned in tin.

There is now real need for civilians to relieve the burden on commercial food sources, transportation, and preservation by growing all food that is practical at home and preserving, storing, and using it over as much of the year as possible.

More than \$200,000,000 worth of vegetables—not counting potatoes and sweetpotatoes—were grown in farm home gardens in 1939. These 4,800,000 home gardens produced vegetables worth a little more than those grown for sale on 3,000,000 acres. Thus, it is obvious that home-grown vegetables can and do furnish a substantial part of our national requirements. If all farmers grow better gardens and all favorably located town dwellers also do their best, a still greater share of the civilian requirements can be met. Now all civilians need to help themselves by home food production insofar as practicable so that more of the country's commercial resources can be devoted to military and lend-lease food requirements.

WE NEED MINERALS AND VITAMINS

Americans, as a group, have not been eating enough of those foods that are rich in the minerals and vitamins necessary for good growth and health. Surveys by nutrition experts and the large number of rejections under the Selective Service Act both emphasize the need for improving our eating habits. Some people have not been eating sufficient quantities of vegetables rich in vitamins and minerals

because they could not get them, but millions more have not eaten enough of these essential vegetables because of lack of knowledge, indifference, or unfortunate food habits, even though they could easily afford and obtain them. National health as well as personal well-being demands that we learn more about what vegetables we need and then make special efforts to use those vegetables effectively. Nutrition experts advise people to get their vitamins from food rather than from indiscriminate use of synthetic preparations.

VEGETABLES AS SOURCES OF MINERALS AND VITAMINS

Vegetables are important foods because of the minerals and vitamins they contain. Their greatest contribution is probably in vitamin A and vitamin C (ascorbic acid), but as a group they also furnish some vitamin B₁ (thiamine), vitamin G (riboflavin), calcium, and iron. Even small amounts of these substances are important, because they supplement what is obtained from other kinds of food.

Vegetables differ greatly in their vitamin and mineral contents. Fortunately, however, some of the commonest and easiest to grow are the most valuable. Table 1 shows which of a number of commonly used vegetables are especially good sources of vitamin A, thiamine, ascorbic acid, riboflavin, calcium, and iron. Others, such as onions and beets, have great practical value in meals because of flavor and color, even though they are not important for good nutrition.

WHO SHOULD GROW VEGETABLES?

Every family living on a farm or country place should, of course, have a vegetable garden. Despite the adverse climate for much of the year in some regions or difficult soil problems, it is practicable at some

TABLE 1.—*Vegetables as sources of vitamins, minerals, and energy*¹

Kind of vegetable	Vitamin A	Thiamine	Ascorbic acid	Riboflavin	Calcium	Iron	Energy value
Beans, lima (green).....		Excellent..		Good.....		Excellent..	Calories pound 595
Beans, snap.....	Excellent..	Good.....	Good.....		Good.....	Good.....	190
Beet greens.....	do.....			Excellent..		Excellent..	150
Cabbage.....		Good.....	Excellent..	Good.....	Good.....		130
Carrots.....	Excellent..	do.....		do.....	do.....		205
Chard.....	do.....				Excellent..	Excellent..	115
Collards.....	do.....	Good.....	Excellent..	Good.....	do.....	Good.....	225
Corn, sweet.....		do.....					510
Kale.....	Excellent..	do.....	Excellent..	Excellent..	Excellent..	Excellent..	225
Lettuce, green.....	do.....	do.....		Good.....	Good.....	Good.....	85
Parsnips.....		do.....	Good.....		do.....		380
Potatoes.....		do.....	do.....				385
Spinach.....	Excellent..	do.....	Excellent..	Good.....		Excellent..	110
Squash, Hubbard.....	do.....						200
Sweet potatoes.....	do.....	Good.....	Good.....				565
Tomatoes, red.....	do.....	do.....	Excellent..				105
Turnip greens.....	do.....	do.....	do.....	Excellent..	Excellent..	Excellent..	165
Turnips, white.....		do.....	do.....		Good.....		155

¹ Data of the Bureau of Home Economics.

season to raise most of the vegetables listed in table 1 with fair success. Even if special handling or treatment, such as irrigation, drainage, or protecting by windbreaks or fences, is necessary, a garden should be grown.

Most people in small towns and villages either have suitable garden spots of their own or can obtain the use of conveniently located small plots of reasonably good soil that are not too steep, too wet, or too shady. In most cases it is not very satisfactory to attempt gardening at any great distance from home. Inconvenience results in neglect. However, small-town and village dwellers who can find good areas near at hand can learn to grow vegetables profitably. Fresh vegetables out of one's own garden give a particular satisfaction and pleasure. Wartime restrictions on travel and on the use of automobiles will keep people at home more and give them more time for gardening. Increased living costs will also doubtless encourage the production of more food at home.

Large-town and city dwellers often do not have suitable space to undertake gardening successfully. Those living in outlying or suburban areas and having large sunny lots, away from interfering buildings, structures, trees, and industrial smoke or gaseous wastes, have a better chance of growing successful gardens than other large-town or city dwellers. It is wasteful, however, to attempt gardening in cramped, poorly drained, poorly lighted spots in the heart of a city or in most highly developed industrial neighborhoods or within the branch or root spread of large trees. If a person insists upon making a garden under such adverse conditions, for exercise or for pleasure or because of the present need, he should realize the odds against profitable yields.

CONSERVE SUPPLIES—FOLLOW THROUGH

No profit will accrue to the Nation or to the individual if prospective gardeners undertake the impossible or even the impracticable. It is wasteful and unwise to devote energy, seeds, fertilizer, and tools to gardening under conditions where success is very doubtful. As long as the United States has the task of helping to feed much of the world, seeds and fertilizer should be carefully conserved. There will probably be enough if they are used with care, but there will be none to waste.

Perhaps the worst waste among gardeners has resulted from neglect and abandonment of gardens planted in a flush of enthusiasm but without adequate means or will to carry each crop through to harvest. The Nation cannot afford such waste of labor and materials when it is at war. Every crop planted should be properly sown at the right time, tended to harvest, then harvested at the proper stage of development, and utilized without waste. Unless the product is actually consumed by those who need it, there is no point in spending seeds, fertilizer, and energy in growing it.

A CONTINUOUS SUPPLY SHOULD BE THE GOAL

When growing vegetables for vitamins and health, a continuous supply of a few health-giving kinds should be the goal. One should guard against planting so much of one vegetable at one time that it will result in a surplus and probable waste, only to be followed by long periods without vegetables. Although some vegetables are suited to planting at intervals over a long period to furnish a continuing supply, others are suited to such a narrow range of changing season that successive plantings are not recommended. Thus, a number of kinds of vegetables or varieties of one kind must be depended upon to provide a continuity of fresh produce. Single plantings, however, of lima beans, some pole snap beans, chard, and tomatoes and, in the South, of fall collards, kale, spinach, and turnips (planted in late summer), remain in a usable stage in the garden for a considerable time after frost.

Some understanding of the behavior of each of these crops under local conditions is necessary in order to plan the best planting schedule and get the most out of the facilities and resources available. Effective gardening requires thought and work before and all through the entire season and, in the South, the year round.

WHAT TO GROW

Primarily, one should grow those vegetables that will be most enjoyed, provided they can be grown successfully in the area. However, if a garden is to contribute to a better fed Nation, it should contain most of the vegetables listed in table 1.

The green leafy vegetables—chard, collards, kale, and turnip greens—are of greatest importance, and one or another of them should be available from early spring onward to severe freezing weather. They should be in every garden and should be eaten frequently.

Tomatoes and beans are so widely adapted that they should be generally and generously grown. Tomatoes are so easily canned that they should be grown for canning as well as for eating fresh.

Beets, carrots, and onions can be stored for some time as well as used fresh; so quantities of these sufficient to afford a supply long after harvest may be planted. Parsnips and salsify can be left in the ground all winter.

Cabbage is easy to grow, but only a few plants per person should be grown for harvesting at one season, as cabbage does not stand long after heading. However, any large surplus of cabbage can be made into sauerkraut, and heads of the Danish type can be stored.

Sweetpotatoes, potatoes, and sweet corn are among the best energy foods, but they require considerable space in the garden. They are not recommended for the small garden, but potatoes might well be included if one has a quarter-acre garden. Sweetpotato vines require so much space that the crop probably should be grown only in the

larger gardens—a third of an acre or more—and then only in regions having a sufficiently long and warm summer. Sweetpotatoes are grown most extensively in the southern half of the United States.

TOOLS

It is always pleasing and sometimes helpful to own a great variety of garden tools—plain and fancy. A large assortment, however, is not necessary, and very simple tools may do quite as good work as fancy or elaborate gadgets.

For a small garden the only essential tools are a spade or spading fork, a steel rake, a common hoe, and a strong cord. Any area should be plowed deeply instead of spaded, if practicable, because spading is very time consuming and laborious. A wheel hoe is a great labor saver, but it is not essential for small gardens. A scuffle hoe, or push hoe, and a trowel are convenient, but not necessary.

If running water is available in regions not normally irrigated, a garden hose is excellent insurance against drought and should be provided—in many regions every householder should have one anyway for protection against incendiary bombs.

During wartime metals and tools are scarce. Buy only your minimum requirements, keep the tools clean and well oiled, and take good care of them.

HOW TO ARRANGE THE PLANTING

If the garden plot slopes appreciably and is subject to washing of the soil, the rows should not run up and down the hill; if the plot is nearly level, the rows should run the long way of the area for convenience in working.

Tall-growing crops should be placed preferably on the north or west side of the garden so that they will not shade the low ones.


Insofar as practicable, the first plantings of small and early vegetables should be along the south or east side, later crops being sown progressively across the area. This orderly procedure helps avoid confusion and damage to the earlier sowings.


In general, flat culture is preferable to and requires less work than growing the crops on raised beds or ridges. Ridges, however, must be used on poorly drained areas where heavy normal rainfall results in frequent surface flooding and on areas where the furrow method of irrigation is to be used.

PLANS FOR GARDENS

It is unwise to recommend any one plan for all regions or for all gardeners. Plans 1 to 3, however, present some simple basic schemes for gardens of different sizes. With a little experience and study of local requirements and possibilities, one can develop improvements that will better adapt the garden to individual needs and localities.


PLAN I. — SCHEME FOR A VERY SMALL GARDEN, 30 BY 50 FEET, APPROXIMATELY 1/30 ACRE!





No one plan suits all gardeners or all parts of the country. This scheme is an example, not a general recommendation

PLAN I. — SCHEME FOR A VERY SMALL GARDEN, 30 BY 50 FEET, APPROXIMATELY $\frac{1}{30}$ ACRE¹



¹ No one plan suits all gardeners or all parts of the country. This scheme is an example, not a general recommendation

Plan 2.—SCHEME FOR A SMALL GARDEN, 35 by 100 feet, approximately $\frac{1}{2}$ acre¹

[Figures at left margin indicate distance in feet between rows. Rows run the long way of the plot]
North or west side of garden

2 feet.	Pole snap beans.	} Kentucky Wonder; $\frac{1}{2}$ pound seed. { After harvesting beans, follow with 3 rows of Purple Top Globe turnips; 1 ounce seed.
2 feet.	Pole snap beans.	
2 feet.	Pole lima beans.	} Carolina or Sieva; $\frac{1}{2}$ pound seed.
2 feet.	Pole lima beans.	
3 feet.	Tomatoes (not staked).	Earliana; $2\frac{1}{2}$ dozen plants; $3\frac{1}{2}$ feet apart.
4 feet.	Tomatoes (not staked).	} Marglobe; 4 dozen plants; 4 feet apart.
4 feet.	Tomatoes (not staked).	
3 feet.	Chard.	Fordhook Giant; 1 ounce seed.
$1\frac{1}{2}$ feet.	Beets.	Crosby Egyptian or Detroit Dark Red; 1 ounce seed. } After harvesting beets, carrots, and turnips, follow these 3 rows August 1 with 2 rows of collards; 200 plants or 2 packets seed.
$1\frac{1}{2}$ feet.	Carrots.	Chantenay or Nantes; $\frac{1}{4}$ ounce seed.
$1\frac{1}{2}$ feet.	Turnips.	Shogoin; $\frac{1}{4}$ ounce seed.
$2\frac{1}{2}$ feet.	Cabbage.	$\frac{1}{2}$ row Golden Acre; $\frac{1}{2}$ row Glory of Enkhuizen; 3 dozen plants of each variety.
$1\frac{1}{2}$ feet.	Onions.	Yellow or Ebenezer; 1 quart sets.
1 foot.	Radishes.	Scarlet Globe; $\frac{1}{3}$ row at 10-day intervals; 1 ounce seed.
1 foot.	Spinach.	Long Standing Bloomsdale; $\frac{1}{2}$ ounce seed. } After harvesting radishes, lettuce, and spinach, follow these 3 rows with 2 rows of U. S. No. 5 Refugee beans; $1\frac{1}{2}$ feet apart; 1 pound seed.
1 foot.	Lettuce.	Black-Seeded Simpson; $\frac{1}{3}$ row at 10-day intervals; 1 packet seed.
$1\frac{1}{2}$ feet.		

¹ No one plan suits all gardeners or all parts of the country. This scheme is an example, not a general recommendation. See p. 2 and bottom of p. 7.

Plan 3.—SCHEME FOR A MEDIUM-SIZE GARDEN spaced for working with horse or small garden tractor, 100 by 150 feet, approximately $\frac{1}{3}$ acre¹

[All rows are 3 feet apart except tomatoes and squash, which are spaced $4\frac{1}{2}$ feet from other crops and $4\frac{1}{2}$ feet apart. Rows run the long way of the plot]

North or west side of garden

Pole snap beans. 2 rows Kentucky Wonder; $\frac{3}{4}$ pound seed. } Follow with 4 rows **turnips**; $1\frac{1}{2}$ ounces seed.

Pole lima beans. 2 rows Carolina or Sieva; $\frac{3}{4}$ pound seed. }

Tomatoes. $\frac{1}{2}$ row Earliana; 2 dozen plants. $\frac{1}{2}$ row Marglobe; 20 plants.

Tomatoes. 2 rows Marglobe; 100 plants.

Squash. $\frac{1}{2}$ row Yellow Bush Scallop; $\frac{1}{2}$ to $\frac{3}{4}$ ounce seed. **Peppers.** $\frac{1}{2}$ row Ruby King; 3 dozen plants.

Parsnips. 1 row Hollow Crown; $\frac{1}{2}$ to $\frac{3}{4}$ ounce seed.

Sweet corn. 4 half rows early; $\frac{3}{4}$ pound seed. 4 half rows late; $\frac{3}{4}$ pound seed. } These rows interplanted with Hubbard squash; $\frac{1}{2}$ row; 1 ounce seed.

Chard. $\frac{1}{2}$ row Fordhook Giant; $\frac{3}{4}$ to 1 ounce seed. New Zealand spinach; $\frac{1}{2}$ ounce seed.

Beets. 1 row Crosby Egyptian; plant half 10 days later; 1 to $1\frac{1}{2}$ ounces seed.

Mustard. $\frac{1}{2}$ row Tendergreen or Giant Curled. **Turnips.** $\frac{1}{2}$ row Shogoin; 1 packet seed each. } Follow with 2 rows of collards; $\frac{1}{2}$ ounce seed or 300 plants.

Turnips. 1 row Shogoin; $\frac{1}{2}$ ounce seed.

Cabbage. $\frac{1}{2}$ row Golden Acre; 4 dozen plants. $\frac{1}{2}$ row Glory of Enkhuizen; 3 dozen plants. } Follow with Detroit Dark Red beets; 3 ounces seed.

Onions. 1 row yellow or Ebenezer; $1\frac{1}{2}$ quarts sets or 500 plants.

Potatoes. 4 rows $\frac{1}{2}$ bushel seed. Follow with U. S. No. 5 Refugee beans; 2 pounds seed.

Spinach. 1 row Long Standing Bloomsdale; $1\frac{1}{2}$ ounces seed.

Carrots. 1 row Chantenay or Nantes; $\frac{1}{4}$ ounce seed.

Peas. 3 rows $1\frac{1}{2}$ pounds seed each of early, medium, and late varieties. Follow with 5 rows of late potatoes; $\frac{1}{2}$ bushel seed.

Radishes. $\frac{1}{2}$ row Scarlet Globe; $\frac{3}{4}$ ounce seed. **Lettuce.** $\frac{1}{2}$ row Curled Simpson; 1 packet seed. } Follow with All Seasons cabbage; 75 plants.

¹ No one plan suits all gardeners or all parts of the country. This scheme is an example, not a general recommendation. See p. 2 and bottom of p. 7.

HOW TO PREPARE AND FERTILIZE THE SOIL

Where the soil is deep, it should be spaded or plowed to a depth of 8 to 10 inches. On thin soils be careful, however, to dig up very little subsoil. Heavy soils should not be worked while wet. Well-rotted leafmold, horse manure or cow manure, or other decayed organic matter, if obtainable, should be worked into the soil in amounts up to about a bushel per 25 square feet (about 20 tons per acre). The spaded-up masses should be crushed and roughly leveled out as the spading progresses.

After the soil of the small garden is spaded, stones or trash at the surface should be removed and clods broken. The clods should be broken up finely with the rake instead of being removed from the garden. Immediately before planting any part of the garden, that part should be finely worked up and smoothed with the rake before laying off the rows.

In locations where the garden spot is not likely to be damaged by washing or blowing during the winter, it is generally helpful to spade or plow it in the fall or early winter and leave it rough. This makes spring preparation easier. If, however, there is danger of washing or blowing, the soil should have a winter cover on it. In the North, gardens may have crops growing in them until so late that it is difficult if not impossible to grow a cover crop of rye or some other suitable kind before winter stops plant growth. In such areas, strawy manure, crop remains, or other coarse plant material may be applied to protect and help hold the soil. This mulching material—or the growing cover—should be turned under in the spring as early as the soil is dry enough to be worked. Very coarse parts of the material that would interfere with the preparation or planting should be removed before spading.

Unless commercial fertilizer is known to be unnecessary for the plot, it should be applied along the row in a band about 3 to 4 inches wide and about 2 inches from the line where the seeds will be sown or the plants set. This can be done by scooping out a wide furrow about 2 inches deep with a good-sized common hoe and then distributing the fertilizer uniformly along the furrow.

Common mixtures like 5-10-5,¹ or those of similar analysis, should be applied at the rate of 1 pound per 30 feet of row (750 pounds per acre) when the rows are 2 feet apart. (An ordinary 10-quart pail holds 15 to 20 pounds of fertilizer.) If the rows are only a foot apart, one-half pound per 30 feet is enough. No more than 1 pound per 30 feet should be applied in bands near the row, regardless of the distance between rows, as otherwise the seedlings may be damaged.

¹ Five percent nitrogen, 10 percent phosphoric acid, and 5 percent potash. If only a 3-8-7 mixture can be obtained, it should be applied at the rate of 1 pound per 24 or 25 feet of row or broadcast and worked in at the rate of 1,200 to 1,500 pounds per acre (3 pounds per 100 square feet, or 10-foot square).

The fertilizer should be mixed thoroughly with the soil and covered about 2 inches deep. The furrow can be partly filled by opening up the row to plant the seed. The fertilizer must not come in contact with the seed. The row for the seed should be opened just before planting. Broadcasting the fertilizer is far easier than applying it in bands on one side and slightly below the seed, but it results in less efficient use of the fertilizer.

HOW TO PLANT

Nearly all gardeners waste seed by sowing it too thickly. This also wastes labor, as the seedlings later must be thinned by hand to a spacing that will allow proper development. Poor growth and poor-quality vegetables are obtained if the seed is sown too thick and the plants are not thinned out to proper spacing in the row. **Don't buy more seed than you need to plant. Don't plant more seed than you need to get a stand.**

Bean and pea seeds should be spaced as the plants are to stand. These vegetables should never be thinned in the rows.

Small seeds like those of carrots, collards, onions, parsnips, spinach, and turnips should be sown three or four times as thick as the plants are to stand finally, since usually many seeds fail to produce good seedlings. Surplus seedlings are thinned out before the plants crowd one another.

Beet and chard "seeds" should be sown no thicker than the plants are to stand, because the "seeds" are really fruits, each containing several seeds. Some thinning is always necessary.

Cabbage, tomato, and onion plants and onion sets should be placed where they are to remain.

Table 2 shows suitable spacing for several crops, the amount of seed required for 1 foot of row and for 100 feet of row, and the proper depth of covering in a good sandy loam. In heavy soils seed should be covered less deeply and in light sandy soils a little more deeply than is indicated.

A mistake many gardeners make is planting the seed too deep. In general, seed should be planted only deep enough to make sure that it is in soil that is moist. Too shallow planting may leave the seed so near the surface that the soil around it dries out before the seed germinates; or the seed may become uncovered by rain. Planting too deep delays germination and may prevent some of the sprouts from pushing through the soil.

In making rows for planting small seeds, about $\frac{1}{2}$ inch deep, the end of the handle of a rake or hoe may be drawn through the soil along the string that marks the row. For such small seeds the soil must be worked until quite fine, smooth, and free from clods, trash, or stones. For larger seeds that are to be planted an inch deep or

TABLE 2.—Seed and space required for certain vegetables when grown in small or intensive gardens

Kind of vegetable	Minimum space between rows	Distance between plants in row	Seed required to plant—			Depth to cover seed
			1 foot of row	100 feet of row	1 acre ¹	
	<i>Inches</i>	<i>Inches</i>	<i>Number</i>		<i>Pounds</i>	<i>Inches</i>
Beans, lima (pole).....	24	24	3-4	¼ pound.....	40-60	1
Beans, snap (pole).....	24	24	3-4	¼ pound.....	60	1
Beets.....	14	2-3	6	1 ounce.....	10-12	1
Cabbage.....	27	15-24		² 50-90.....	² 15, 000	
Carrots.....	14	2-3	20-25 ³	¼ ounce.....	4	½
Chard.....	18	4-6	3-4	1 ounce.....	8-10	1
Collards.....	18	15-18	³ 3-4	1 packet.....	³ ¼	½
Corn, sweet.....	36	15	³ 3-4	¼ pound.....	15	1
Kale.....	18	12	³ 3-4	1 packet.....	4	½
Lettuce.....	15	12	8-10 ⁴	1 packet.....	2	½
Mustard.....	15	4-6	8-10	¼ ounce.....	2	½
Onions.....	14	2-3	15-20	1 quart ⁴		½
Parsnips.....	18	2-3	15-20	½ ounce.....	3	½
Peas.....	18	1	12-15	1 pound.....	150	1-1½
Potatoes.....	24	12		6-8 pounds.....	900	4
Radishes.....	12	2	12-15	1 ounce.....	12	½
Spinach.....	12	3-4	10-15	½ ounce.....	12	½
Squash, Hubbard.....	100	30	³ 4-5	1 ounce.....	2	1-1½
Sweetpotatoes.....	36	15		² 80.....	² 12, 000	
Tomatoes (staked).....	36	24		² 50.....	² 7, 500	
Turnips.....	14	2-3	20-25	¼ ounce.....	2	½

¹ Average instead of very intensive rate of planting for use in calculating seed requirements for larger gardens.

² Plants.

³ Several seeds planted in one spot where the plants are to stand.

⁴ Sets.

deeper, the row is best opened up with the corner of a common hoe. With a little practice it will be possible to make the trench of uniform and proper depth for the kind of seed to be planted. Irregular depth or irregular covering often causes irregular "come-up" and stand of plants.

TIME TO PLANT DIFFERENT VEGETABLES

Because of the great diversity of climates and seasons over the country, no detailed information on planting dates can be given in this publication. Vegetables, however, may be roughly classified and sown according to their hardiness and temperature requirements. Gardeners should consult their experienced neighbors and local agricultural advisers or obtain the more detailed publications available on gardening (p. 2). A rough timetable of planting is shown as table 3.

TABLE 3.—Approximate time to plant certain vegetables

Early-spring plantings		Late-spring or summer plantings		Late-summer or fall plantings (6 to 8 weeks before fall freeze)
4 to 6 weeks before frost-free date	2 to 4 weeks before frost-free date	Frost-free date	2 to 6 weeks after frost-free date	
Cabbage plants.	Beets.	Beans.	Beans, snap.	Beets.
Lettuce.	Carrots.	Beets.	Beets.	Collards.
Onions.	Chard.	Corn, sweet.	Corn, sweet.	Kale.
Peas.	Lettuce.	Squash.	Sweetpotatoes.	Mustard.
Potatoes.	Mustard.	Tomato plants.		Spinach.
Spinach.	Peas.			Turnips.
Turnips.	Parsnips.			
	Radishes.			

CULTIVATING AND MULCHING

All weeds must be kept under control by thorough shallow cultivation or hoeing. Vegetable crops should not be cultivated deeply because of danger to the roots that grow near the surface. Weeds that take root again readily after hoeing or pulling should be carried out of the garden. The garden should be cultivated as soon as the soil is dry enough after each rain or irrigation and as often in addition as is necessary to keep the weeds down. There is no proved benefit from stirring an already cultivated soil that is free of weeds.

Care should be taken to avoid trampling and packing the soil so far as practicable. Mulching between the rows with straw, dried lawn clippings, leaves, or similar material will help conserve moisture and keep down weeds.

WATERING OR IRRIGATION

Frequent light sprinkling or irrigation is a bad practice. If water is to be applied, the garden should be thoroughly and deeply soaked, as by a fairly heavy rain, and watered again only when the soil shows signs of becoming dry.

PROPER HARVESTING AND USE OF PRODUCTS

Too many growers, in attempts to get larger growth and yield, delay harvest beyond the stage of best quality. No vegetable should be allowed to become tough, coarse, overgrown, and unpalatable before being harvested. Quantity is important, but so is quality. Large size in a product is, of itself, of little value. Indeed, excessive size is generally associated with mediocre, if not low, quality.

The sooner vegetables can be used after harvest, the better. If they must be kept awhile, they should generally be kept in a cool moist place.

Most people cook vegetables too long; for example, cabbage and snap beans are often overcooked. This destroys much of the vitamin content. They also cook them in too much water, which removes and wastes part of the valuable minerals. Soda should never be added to green vegetables to set the color, as it destroys vitamin C.

HOME STORAGE

Many of the vegetables referred to in this bulletin can be stored for later use. For successful storage, however, they must be fully grown or mature and free from cuts, bruises, and injuries from insects or diseases. If the gardener can properly produce more of these storable items than needed during the growing season, they should be grown and put away carefully for later use.

Surplus beans and peas may be allowed to ripen on the plants. The pods may then be hand-picked, threshed, dried, and stored in closed containers to keep insects out. The seeds are easily threshed by placing the pods in large cloth bags, drying thoroughly, then pounding, and shaking vigorously.

Fall carrots, beets, turnips, and potatoes can be kept in a cool, moist cellar but should not freeze. Parsnips and salsify may be either left in the soil all winter or stored like other root crops. Sweetpotatoes and winter squashes require warmer storage than the root crops do: For sweetpotatoes, a temperature of 55° to 60° F. and a medium relative humidity (75 to 80 percent); and for winter squashes, 50° to 60° and 70 to 75 percent relative humidity.

Onions require cool, dry storage conditions, as in an attic or shed where they can be protected from freezing, preferably about 35° F.

For further information on storage, write Office of Information, United States Department of Agriculture, Washington, D. C.

DEHYDRATION AND CANNING

Because of the shortage of metals, especially tin, and the difficulties of storage in the natural state, there is a greatly renewed interest in the preservation of vegetables by dehydration or drying. As mentioned in the preceding section, ripe pea and bean seed dry naturally and keep well if insects and moisture are kept away from them. Cabbage, onions, carrots, and sweetpotatoes are being dehydrated commercially on a large scale, as are a number of other vegetables. Sweet corn in the roasting-ear stage has been successfully dried for generations by thrifty farm wives.

It is not the purpose of this bulletin to go into the details of food preservation. However, the possibilities of drying and canning should not be overlooked by those gardeners who can thus add to their food supplies. The following publications on these subjects can be obtained free from the Office of Information, United States Department of Agriculture, Washington, D. C.:

Farmers' Bulletin 1918, *Drying Foods for Victory Meals.*

Farmers' Bulletin 1762, *Home Canning of Fruits, Vegetables, and Meats.*

